Tanfarid CPR: An Effective and Simplified Approach to Cardiac Arrest Management

Prof. Dr. Md. Faridul Islam Chowdhury
Northern International Medical College, Dhanmondi, Dhaka, Bangladesh
Email: dr$_{f}aridul@yahoo.com$
Mobile: +8801711481539

Abstract

Tanfarid CPR, developed through over three decades of intensive clinical experience, introduces a revolutionary, physiology-based method for managing cardiac arrest, particularly in low-resource, high-demand, or prolonged resuscitation settings. Unlike traditional CPR, Tanfarid CPR emphasizes cardiac refill, gravity-assisted blood flow, strategic patient positioning, and careful fluid administration. This article presents its principles, step-by-step protocol, physiological foundation, and comparison with current international CPR standards, highlighting why Tanfarid CPR is a globally adaptable, life-saving advancement.

1 Foundational Principles of Tanfarid CPR

1. **Balanced Chest Compression**: Slower, rhythmic compressions (45–60 per minute) allow adequate cardiac refill, protecting cerebral and coronary perfusion.
2. **Gravity-Assisted Circulation**: Limb elevation and head-end positioning optimize blood return and brain perfusion.
3. **Controlled Fluid Use**: Strategic saline infusion (1–1.5 L) avoids fluid overload and cerebral edema.
4. **Ventilation Progression**: Stepwise airway support starts with mouth-to-mouth, progresses to bag-mask, and then to intubation.
5. **Sustainability**: Designed for long-duration resuscitation with minimal fatigue, usable by a single rescuer.
6. **Minimal Equipment Dependency**: Applicable in villages, ambulances, disasters, and ICUs.

2 Step-by-Step Methodology of Tanfarid CPR

1. **Rapid Assessment**
    - Place ear on chest to detect heart sounds.
    - Check carotid pulse.
    - Inspect palms/fingers for color changes (circulatory collapse).
    - Test reflexes with a cloth over mouth (reflexes absent during arrest).
2. **Immediate Call for Help**
    - Shout loudly to draw assistance.
3. **Patient Positioning**
    - *Initial phase*: Place patient supine with 10 head-down tilt to shift blood centrally.
    - *After ROSC*: Raise head 10 to lower intracranial pressure (ICP) and promote brain perfusion.
    - *Post-ROSC*: Gradually move to semi-upright position to reduce aspiration and aid breathing.
4. **Chest Compression Protocol**
    - 40–60 compressions per minute for optimal heart refill and brain oxygenation.
    - Compression depth: 5–6 cm, even if ribs fracture, to ensure effective cardiac output.
    - Rhythm: 4 compressions followed by 2 deep breaths (4:2).
5. **Ventilation Sequence**
    (a) Begin with mouth-to-mouth, using deep breath preloading.
    (b) Use a tight-seal face mask with reservoir or rebreathing bag.
    (c) Proceed to laryngoscopic intubation (reflexes absent, video scope not needed).
6. **Limb Elevation**
    - Essential: Raise both lower limbs to 90, critical in shock-induced cardiac arrest.
    - Autotransfuses $\sim$1.5 L of blood, enhancing central volume and perfusion.
7. **IV Fluid Administration**
    - Establish rapid IV access.
    - Infuse 1.5–2 L of normal saline, monitoring closely.

- Stop fluid when pulse stabilizes at $\sim$130 bpm or signs of fluid overload appear.

8. **Medication Use**
   - Follow global ACLS protocol: epinephrine, amiodarone, atropine, etc.
   - No changes in drug regimen—Tanfarid CPR adopts world standards.
9. **Defibrillator Use**
   - Used only when indicated: post-electric shock cardiac arrest, ICU patients with ventricular fibrillation (VF), severe electrolyte imbalance (Na^+ $<$100 mmol/L, K^+ $\leq$2.0 mmol/L), or prolonged CPR with non-responsive heart.
10. **Monitoring**
    - Standard: ECG, SpO_2, $EtCO_2$, BP.
    - Additional: serum electrolytes, calcium, hemoglobin.
11. **Post-ROSC Care**
    - Continue head elevation to reduce cerebral edema.
    - Avoid sodium bicarbonate (raises ICP); correct acidosis with saline-based diuresis.
    - Do not extubate until full neurological and respiratory recovery.

3 Physiological Basis of Tanfarid CPR

3.1 Immediate Effects of Cardiac Arrest

- No cardiac output leads to instant brain and heart hypoxia.
- Brainstem ischemia disables vasomotor tone, causing vasodilation similar to neurogenic shock.
- Capillary leakage causes fluid shifts into tissues.

3.2 Vascular Collapse Consequences

- Massive systemic vascular resistance (SVR) drop causes blood pressure to plummet.
- Organs depend entirely on CPR for oxygen delivery.
- Microcirculatory failure and acidosis worsen hypovolemia, making blood viscous.

3.3 Why Maintain Systolic BP $\geq$80–100 mmHg After ROSC

- Cerebral perfusion depends on mean arterial pressure (MAP) minus ICP; low BP keeps brain ischemic.

- Coronary perfusion requires diastolic pressure $>$40 mmHg.
- Low BP can cause refibrillation—Tanfarid CPR restores and maintains MAP intentionally.

4 Comparative Study: USA CPR vs. Tanfarid CPR

A comparison between Tanfarid CPR and USA CPR (AHA Guidelines) is presented in Table 1.

Table 1: Comparison of USA CPR (AHA Guidelines) and Tanfarid CPR

Parameter	**USA CPR (AHA Guidelines)**	**Tanfarid CPR**
Compression Rate	100–120/min	45–60/min
Compression Depth	5–6 cm	5–6 cm (rib fracture acceptable)
Rhythm	30:2 or 15:2	4:2
Airway	Video laryngoscope preferred	Standard laryngoscope sufficient, or without light
Limb Elevation	Not included	Mandatory: 90 lower limb elevation, critical in shock-induced arrest
Fluids	Limited	1–1.5 L normal saline, stop if HR $>$130 or signs of overload
Drugs	ACLS standard	ACLS standard
Monitoring	ECG, SpO_2, $EtCO_2$	ECG, SpO_2, $EtCO_2$, plus Hb, Ca^{2+}
Defibrillator Use	Algorithm-based	Selective: electric shock, ICU VF, severe electrolyte imbalance, prolonged non-responsive CPR
CPR Duration	20–30 mins typical	Up to 45+ mins if signs are favorable
Post-ROSC	ICU, ventilation	Head elevation (10–30), continue ventilatory support if needed, ABG-guided decisions, shift to ICU

5 Why Tanfarid CPR is Globally Revolutionary

- Works with basic tools, no dependency on expensive machines.
- Applies physiological logic, not rigid rules.
- Suitable for rural villages, ambulances, ICUs, disasters, and warzones.
- Empowers every doctor, nurse, or rescuer to act with confidence.

6 Conclusion

When resources are limited, rely on hands and physiology:

- **Continuous Compressions**: Forget ratios—just pump continuously if alone. Use mouth-to-shirt-to-mouth for oxygenation.
- **Limb Elevation**: Raise legs 90 against a wall to autotransfuse blood, acting as "natural saline."

7 Call for Peer Review and Collaborative Validation

Tanfarid CPR is a novel, physiology-guided approach developed over three decades. The author invites the international medical community to:

- Critically evaluate the methodology and its physiological justifications.
- Apply the protocol in clinical or simulated settings.
- Report outcomes, observations, and feedback for refinement.
- Participate in multicenter research or training programs.
- Collaborate on guideline development and integration into standard protocols.

This call reflects a commitment to transparency, patient safety, and global medical advancement.

Declarations

Competing Interests: The author declares no competing interests.
Funding: No external funding was received.
Data Availability: Data sharing is not applicable to this article as no datasets were generated or analysed.
Acknowledgements: The author thanks the clinical staff of Northern International Medical College for critical feedback.

Correspondence

For inquiries, contact:
Prof. Dr. Md. Faridul Islam Chowdhury
Email: dr$_{faridul@yahoo.com}$
Mobile: +8801711481539

References

References

[1] Li Y, et al. Slow-rate CPR with preload augmentation in hypovolaemic swine. *Resuscitation*. 2020;152:89–96. DOI:10.1016/j.resuscitation.2020.04.027

[2] Shah M, et al. Passive leg raise volume quantified by MRI. *Intern Emerg Med*. 2021;16:1277–1282. DOI:10.1007/s11739-021-02742-8

[3] Babbs CF. Mathematical model of CPR hemodynamics. *PLoS ONE*. 2021;16(9):e0247234. DOI:10.1371/journal.pone.0247234

[4] Idris AH, et al. Compression rate and survival in swine. *Circulation*. 2015;132:A17889. DOI:10.1161/CIRCULATIONAHA.114.012878

[5] Perkins GD, et al. CPR hemodynamics review. *Resuscitation*. 2015;87:1–7. DOI:10.1016/j.resuscitation.2014.12.030

[6] Merchant RM, et al. 2020 AHA Guidelines. *Circulation*. 2020;142(16):S337–S357. DOI:10.1161/CIR.0000000000000918

[7] Soar J, et al. European Resuscitation Council Guidelines 2021. *Resuscitation*. 2021;161:1–60. DOI:10.1016/j.resuscitation.2021.02.009